













- , — (1998): Contribution to the crystal chemistry of thallium sulfosalts. IV. Modular description of Tl–As–Sb sulfosalts rebulite and jankovcité. *N. Jb. Miner. Abh.*, 174, 181-210.
- , — (1999): Gillyuite  $Tl_2(As, Sb)_3S_{13}$ : Reinterpretation of the crystal structure and order-disorder phenomena. *Am. Mineral.*, 84, 400-406.
- Makovicky, E. & Hyde, B.G. (1981): Non-commensurate (misfit) layer structures. *Structure and Bonding*, 46, 101-170.
- , — (1992): Incommensurate, two-layer structures with complex crystal chemistry: Minerals and related synthetic materials. *Sci. Forum*, 100, 101, 1-100.
- Makovicky, E. & Karup-Møller, S. (1977a): Chemistry and crystallography of the illianite homologous series. Part I: General properties and definitions. *N. Jb. Miner. Abh.*, 130, 264-287.
- , — (1977b): Chemistry and crystallography of the illianite homologous series. Part II: Definition of new minerals: eskimoite, vikingite, ourayite and treasurite. Redefinition of schirmerite and new data on the illianite–gustavite solid solution series. *N. Jb. Miner. Abh.*, 131, 56-82.
- , — (1984): Ourayite from Ivigtut, Greenland. *Can. Mineral.*, 22, 565-575.
- , — (1986): New data on giessenite from the Bjørkåsen sulfide deposit at Otdalen, Northern Norway. *Can. Mineral.*, 24, 21-25.
- Makovicky, E. & Makovicky, M. (1978): Representation of compositions in the bismuthinite–aikinite series. *Can. Mineral.*, 16, 405-409.
- Makovicky, E. & Mumme, W.G. (1979): The crystal structure of benjaminite  $Cu_{50}Pb_{40}Ag_{2.30}Bi_{6.80}S_{12}$ . *Can. Mineral.*, 17, 607-618.
- , — (1983): The crystal structure of ramdohrite,  $Pb_{11}Ag_3S_{24}$ , and its implications for the andorite group and zinckenite. *N. Jb. Miner. Abh.*, 147, 58-79.
- , — (1984): The crystal structures of izoklakeite, dadsonite and jaskólskiite. 13th Int. Congress of Crystallogr., Hamburg, 9-18 August, Collected Abstracts, C-246.
- , — (1986): The crystal structure of isoklakeite,  $Pb_{51.3}Sb_{20.4}Bi_{19.5}Ag_{12}Cu_{2.9}Fe_{0.7}S_{114}$ . The kobellite homologous series and its derivatives. *N. Jb. Miner. Abh.*, 153, 121-145.
- Makovicky, E. & Nørrestam, R. (1985): The crystal structure of jaskólskiite,  $CuPb_{2+x}(Sb, Bi)_{25-x}S_5$  ( $x = 0.2$ ), a member of the meneghinite homologous series. *Z. Kristallogr.*, 171, 179-194.
- Makovicky, E. & Skinner, B. (1975): Studies of the sulfosalts of copper. IV. Structure and twinning of sinnerite,  $As_4S_9$ . *Am. Mineral.*, 60, 998-1012.
- , — (1979): Studies of the sulfosalts of copper. VII. Crystal structures of the exsolution products  $Cu_9Sb_4S_{13}$  and  $Cu_{3.8}Sb_4S_{13}$  of unsubstituted synthetic tetrahedrite. *Can. Mineral.*, 17, 619-634.
- Makovicky, E., Mumme, W.G., Watts, J.A. (1977): The crystal structure of synthetic pavonite,  $Ag_5S_8$ , and the definition of the pavonite homologous series. *Can. Mineral.*, 15, 339-348.
- Makovicky, E., Johan, Z., Karup-Møller, S. (1980): New data on bukovite, thalcusite, chalcothallite and rohaltite. *N. Jb. Miner. Abh.*, 138, 122-146.
- Makovicky, E., Mumme, W.G., Hoskins, B.F. (1991): The crystal structure of Ag–Bi-bearing heyrovskýite. *Can. Mineral.*, 29, 553-559.
- Makovicky, E., Mumme, W.G., Madsen, I.C. (1992): The crystal structure of vikingite. *N. Jb. Miner. Mh.*, 1992(10), 454-468.
- Makovicky, E., Leonardsen, E., Moëlo, Y. (1994): The crystallography of lengenbachite, a mineral with the non-commensurate layer structure. *N. Jb. Miner. Abh.*, 2, 169-191.
- Makovicky, E., Balč-Zunič, T., Topa, D. (2001a): The crystal structure of neyite,  $AgCu_6Pb_{25}Bi_{26}S_{68}$ . *Can. Mineral.*, 39, 1365-1376.
- Makovicky, E., Topa, D., Balč-Zunič, T. (2001b): The crystal structure of paarite, the newly discovered 56 Å derivative of the bismuthinite–aikinite solid solution series. *Can. Mineral.*, 39, 1377-1382.
- Makovicky, E., Sørensen, I., Karup-Møller, S. (2002): The crystal structure of  $Cu_4Bi_4S_8$ . *Z. Kristallogr.*, 217, 597-604.
- Makovicky, E., Balč-Zunič, T., Karanović, L., Poleti, D., Peck, J. (2004): Structure refinement of natural robinsonite,  $Sb_4S_{13}$ : cation distribution and modular analysis. *N. Jb. Miner. Mh.*, 2004(2), 49-67.
- Makovicky, E., Karanović, L., Poleti, D., Balč-Zunič, T., Paar, W.H. (2005): Crystal structure of copper-rich unsubstituted tennantite,  $Cu_{25}As_4S_{13}$ . *Can. Mineral.*, 43, 679-688.
- Makovicky, E., Balč-Zunič, T., Karanović, L., Poleti, D. (2006a): The crystal structure of kirkiite,  $Pb_6Bi_3As_3S_{19}$ . *Can. Mineral.*, 44, 177-188.
- Makovicky, E., Topa, D., Mumme, W.G. (2006b): The crystal structure of dadsonite. *Can. Mineral.*, 44, 1499-1512.
- Mandarino, J.A. & Back, M.E. (2004): Fleischer's glossary of mineral species 2004. The Mineralogical Record Inc., Tucson, 309 p.
- Mantienne, J. (1974): La minéralisation thallifère de Jas-Roux (Hautes-Alpes). Thesis, University of Paris, 153 p.
- Martin, R.F. (2003): Encyclopedia of mineral names: third update. *Can. Mineral.*, 41, 1075-1096.
- Martin, R.F. & Blackburn, W.H. (1999): Encyclopedia of mineral names: first update. *Can. Mineral.*, 37, 1045-1078.
- , — (2001): Encyclopedia of mineral names: second update. *Can. Mineral.*, 39, 1199-1218.
- Marumo, F. (1967): The crystal structure of nowackiite,  $Cu_6Zn_3As_4S_{12}$ . *Z. Kristallogr.*, 124, 351-368.
- Marumo, F. & Nowacki, W. (1965): The crystal structure of rathite-I. *Z. Kristallogr.*, 122, 433-456.
- , — (1967a): The crystal structure of hatchite  $PbTiAg_3S_5$ . *Z. Kristallogr.*, 125, 249-265.
- , — (1967b): The crystal structure of dufrénoysite,  $Pb_5S_4O$ . *Z. Kristallogr.*, 124, 409-419.
- Matsumoto, T. & Nowacki, W. (1969): The crystal structure of trechmannite,  $AgAs_5$ . *Z. Kristallogr.*, 129, 163-177.
- Matsushita, Y. & Takéuchi, Y. (1994): Refinement of the crystal structure of hutchinsonite,  $TlPb_3S_5$ . *Z. Kristallogr.*, 209, 475-478.
- Matsushita, Y. & Ueda, Y. (2003): Structure and physical properties of 1D magnetic chalcogenide, jamesonite ( $F_6Pb_5S_{14}$ ). *Inorg. Chem.*, 42, 7830-7838.
- Matsushita, Y., Takéuchi, Y., Nishi, F. (1997): The crystal structure of semseyite,  $Pb_5S_{21}$ . In "Tropochemical cell-twinning: A Structure-Building Mechanism in Crystalline Solids". Terra Scientia Publishing Company, Tokyo, 288-293.
- Matzat, E. (1979): Cannizzarite. *Acta Cryst.*, B35, 133-136.
- McQueen, K.G. (1987): A second occurrence of falkmanite: Pinnacles mine, Broken Hill, New South Wales. *Can. Mineral.*, 25, 15-19.
- Meerschaut, A., Palvadeau, P., Moëlo, Y., Orlandi, P. (2001): Lead-antimony sulfosalts from Tuscany (Italy). IV. Crystal structure







## Part II. Review of sulfosalt systematics

IMA-COM Sulfosalt Sub-Committee

### Introduction: general presentation of sulfosalt species

This general presentation takes into account the sulfosalt species given by “Fleischer’s Glossary of Mineral Species” (Mandarino & Back, 2004) (see also Blackburn & Dennen, 1997; Martin & Blackburn, 1999, 2001; Martin, 2003), plus the new species published or approved recently by the CNMNC-IMA (see its website). The presentation is concerned with more than 220 sulfosalt species, for which an alphabetical list is given at the end of the text, together with an appendix that lists discredited species.

As the crystal-chemical classification of sulfosalts is incomplete at present, the following general presentation of sulfosalt mineral species is subdivided into large chemical groups. Within each group, subdivisions are generally based on well-defined structure types.

Sulfosalt species whose specific crystal structure does not have a close relationship to those of other species are indicated separately as “*Single type*”. If the crystal structure of a species is not known, this species is classified, as much as possible, with sulfosalts that have a similar chemistry.

#### About the references

To reduce as much as possible the number of references cited in this review, only the following have been included:

- systematically, the studies presenting the crystal structures of the sulfosalt species (noted “STR” afterwards), but also taking into account data obtained on synthetic compounds (“synth.” afterwards);
- recent papers that define sulfosalt species (since 1990, or older, when necessary);
- all references needed for the presentation and discussion of problems of definition and nomenclature.

Crystallographic data (unit-cell parameters, symmetry, space group) have been avoided, except when a change in symmetry or space group appears crucial for the distinction between two very close species (*e.g.*, giessenite *versus* izoklakeite).

All other references and basic data are available in fundamental books on systematic mineralogy (*e.g.*, Strunz & Nickel, 2001; Mandarino & Back, 2004), as well as in PDF (JCPDF) or ICSD (FIZ – Karlsruhe) databases. Concerning the crystal structures, especially noteworthy is the extensive work of Dr Y. Matsushita, who has compiled systematically all chalcogenide and related structures, both of natural and synthetic phases. Access to the data-library is free at <http://www.crystalmaker.co.uk/library/chalcogenides.html>.

Where problems are present regarding the definition of a species, relevant comments are given after the presentation of each species or group. The aim is to present the current status of sulfosalt definition, nomenclature and classification for all specialists interested in this field of research,

thereby pointing out various unsolved questions and facilitating the discovery of new mineral species.

### 1. Sulfosalts with atom ratio of cation/chalcogen = 1

#### 1.1. Binary sulfosalts ( $MPnCh_2$ ), where $M$ = univalent cation (Cu, Ag, Tl); $Pn$ = pnictogen (As, Sb, Bi); $Ch$ = chalcogen

These sulfosalts are presented according to the organisation of pnictogen polyhedra.

##### 1. Matildite isotypic series (trigonal derivatives of $PbS$ , according to $(PbS)_{111}$ slices)

###### **Matildite**, $AgBiS_2$

STR (synth.): Geller & Wernick (1959).

###### **Bohdanowiczite**, $AgBiSe_2$

STR (synth.): Geller & Wernick (1959).

###### **Volynskite**, $AgBiTe_2$

STR (synth.): Pinsker & Imamov (1964).

All these structures could also be considered as derivatives of the  $CdI_2$  archetype (single layer of  $BiCh_6$  octahedra), with Ag atoms intercalated between the layers (so-called “intercalation compounds”). However, these old structure determinations appear to be (pseudo)cubic approximations, as it is unrealistic to consider regular  $BiCh_6$  octahedra because of the lone-electron-pair of  $Bi^{3+}$ .

##### 2. Aramayoite isotypes

###### **Aramayoite**, $Ag_3Sb_2(Bi, Sb)S_6$

###### **Baumstarkite**, $Ag_3Sb_3S_6$

Definition of baumstarkite and STR of aramayoite and baumstarkite are given by Effenberger *et al.* (2002).

##### 3. (*Single type*)

###### **Cuboargyrite**, $AgSbS_2$

Defined by Walenta (1998).

STR (synth.): Geller & Wernick (1959).

##### 4. (*Single type*) (*sheared derivative of $SnS$ archetype*)

###### **Miargyrite**, $AgSbS_2$

STR: Smith *et al.* (1997).

##### 5. (*Single type*)

###### **Smithite**, $AgAsS_2$

STR: Hellner & Burzlaff (1964). In the structure, As in triangular pyramidal coordination forms  $As_3S_6$  trimers arranged in columns parallel to  $b$ .

















- of pillaitite,  $Pb_3Sb_{10}S_{23}ClO_{0.5}$ , an expanded monoclinic derivative of hexagonal  $Bi(Bi_2S_3)_9I_3$ . *Eur. J. Mineral.*, 13, 779-790.
- Megarskaya, L., Rykl, D., Taborsky, Z. (1986): Zoubekite,  $AgPb_4Sb_4S_{10}$ , a new mineral from Příbram, Czechoslovakia. *Jb. Miner. Mh.*, 1986(1), 1-7.
- Meisser, N., Schenk, K., Berlepsch, P., Brugger, J., Bonin, M., Criddle, A.C., Thélin, P., Bussy, F. (2007): Pizgrischite,  $(Cu, Fe)Cu_4PbBi_7S_{35}$ , a new sulfosalt from the Swiss Alps: description, crystal structure and occurrence. *Can. Mineral.*, 45, 1229-1245.
- Mellini, M. & Merlino, S. (1979) Versiliaite and apuanite: Derivative structures related to schafarzikite. *Am. Mineral.*, 64, 1235-1242.
- Miehe, G. (1971): Crystal structure of kobellite. *Nature, Phys. Sci.*, 231, 133-134.
- Moëlo, Y. (1989): Antimoine dans la nueldite associée à de la friedrichite (commune des Houches, Alpes de Haute-Savoie); redétermination cristallochimique de la nueldite. *C. R. Acad. Sci. Paris*, 309, Sér. II, 1659-1664.
- Moëlo, Y., Borodaev, Yu.S., Mozgova, N.N. (1983): Association twinnite-zinkénite-plagionite du gisement complexe à Sb-Pb-Zn de Rujevac (Yougoslavie). *Bull. Minéral.*, 106, 505-510.
- Moëlo, Y., Makovicky, E., Karup-Møller, S. (1984a): New data on the minerals of the andorite series. *Jb. Miner. Mh.*, 1984(4), 175-182.
- Moëlo, Y., Mozgova, N., Picot, P., Bortnikov, N., Vrublevskaya, Z. (1984b): Cristallographie de l'owyheeite: nouvelles données. *Tschermaks Mineral. Petrogr. Mitt.*, 32, 271-284.
- Moëlo, Y., Jambor, J.L., Harris, D.C. (1984c): Tintinaite et sulfosels associés de Tintina (Yukon): la cristallographie de la série de la kobellite. *Can. Mineral.*, 22, 219-226.
- Moëlo, Y., Makovicky, E., Karup-Møller, S. (1989): Sulfures complexes plombo-argentifères : minéralogie et cristallographie de la série andorite-zélyite,  $(Pb, Mn, Fe, Cd, Sn)$   $(Ag, Cu)_x(Sb, Bi, As)_{2-x}(S, Se)_y$ . Documents du BRGM 167, BRGM éd., Orléans, 107 p.
- Moëlo, Y., Makovicky, E., Karup-Møller, S., Cerville, B., Maurel, C. (1990): La lévyclaudeite,  $Pb_5Sn_7Cu_8(Bi, Sb)_8S_{28}$ , une nouvelle espèce minérale à structure incommensurable, de la série de la cylindrite. *Eur. J. Mineral.*, 2, 711-723.
- Moëlo, Y., Roger, G., Maurel-Palacin, D., Marcoux, E., Laroussi, A. (1995): Chemistry of Pb-(Cu, Fe)-(Sb, Bi)-sulfosalts from France and Portugal, and cationic substitutions in the Cu-poor part of the  $Pb_2S_2-Cu_2S-Sb_2S_3-Bi_2S_3$  system. *Mineral. Petrol.*, 53, 229-250.
- Moëlo, Y., Meerschaut, A., Makovicky, E. (1997): Re nément of the crystal structure of nueldite,  $Pb_2Cu_{1.4}(Pb_{0.4}Bi_{0.4}Sb_{0.2})Bi_2S_7$ : structural relationships and genesis of complex lead sulfosalt structures. *Can. Mineral.*, 35, 1497-1508.
- Moëlo, Y., Meerschaut, A., Orlandi, P., Palvadeau, P. (2000): Lead-antimony sulfosalts from Tuscany (Italy): II - Crystal structure of scainiite,  $Pb_4Sb_{30}S_{54}O_5$ , an expanded monoclinic derivative of  $Ba_{12}Bi_{24}S_{48}$  hexagonal sub-type (zinkenite series). *Eur. J. Mineral.*, 12, 835-846.
- Moëlo, Y., Palvadeau, P., Meisser, N., Meerschaut, A. (2002): Structure cristalline d'une ménéghinite naturelle pauvre en cuivre,  $Cu_{0.58}Pb_{1.72}(Sb_{0.74}Bi_{0.24})S_{24}$ . *C. R. Geoscience*, 334, 529-536.
- Moh, G. (1987): Mutual Pb/Sr<sup>2+</sup> substitution in sulfosalts. *Mineral. Petrol.*, 36, 191-204.
- Mozgova, N.N. (1984): Principles of classification of sulfosalts. in Proceedings of the 27th International Geological Congress, Moscow, August 4-14, Section 10, 53-65.
- (2000): Sulfosalt mineralogy today. IMA COM short course "Modern approaches to ore and environmental mineralogy", Espoo, Finland, June 11-17, 4 p. (extended abstract).
- Mozgova, N.N. & Borodaev, Yu.S. (1972): Homologies within the semseyite-plagionite series. *Zapiski Vsesoyuz. Mineral. Obshch.*, 3, 299-312 (in Russian).
- Mozgova, N.N. & Bortnikov, N.S. (1980): Non-stoichiometry of acicular lead sulfosalts. *Geokh. Mineral.*, XXVth Intern. Geol. Congr., Moscow, 126-138 (in Russian).
- Mozgova, N.N., Organova, N.I., Gorshkov, A.I. (1976): Structural resemblance between incaite and frankite. *Dokl. Acad. Nauk SSSR*, 228, 110-113 (in Russian).
- Mozgova, N.N., Bortnikov, N.S., Tsepin, A.I., Borodaev, Yu.S., Vrublevskaya, S.V., Vyalsov, L.N., Kuzmina, O.V., Sivtsov, A.V. (1983): Falkmanite,  $Pb_4Sb_{36}S_{11}$ , new data and relationship with sulphantimonites of lead (re-examination of type material from Bayerland Mine, Bavaria). *Jb. Miner. Abh.*, 147, 80-98.
- Mozgova, N.N., Vrublevskaya, Z.V., Sivtsov, A.V. (1984): New data on the boulangerite homologous series. *Dokl. Acad. Nauk SSSR*, 274, 169-172 (in Russian).
- Mozgova, N.N., Organova, N.I., Borodaev, Yu.S., Ryabeva, E.G., Sivtsov, A.V., Getmanskaya, T.I., Kuzmina, O.V. (1988): New data on cannizzarite and bursaitite. *Jb. Miner. Abh.*, 158, 293-309.
- Mozgova, N.N., E mov, A.V., Nenasheva, S.N., Golovanova, T.I., Sivtsov, A.V., Tsepin, A.I., Dobretsova, I.G. (1989): New data on diaphorite and brongniardite. *Zapiski Vsesoyuz. Mineral. Obshch.*, 5, 47-63 (in Russian).
- Mozgova, N.N., Nenasheva, S.N., Chistyakova, N.I., Mogilevkin, S.B., Sivtsov, A.V. (1990): Compositional fields of minerals in the bismuthinite-aikinite series. *Jb. Miner. Mh.*, 1990(1), 35-45.
- Mozgova, N.N., Moëlo, Y., Borodaev, Yu.S., Nenasheva, S.N., E mov, A.V. (1992): Wittite with Se-rich cosalite and bismuthinite from Nevskoe tin deposit (Magadan district, Russia). *Mineral. Petrol.*, 46, 137-153.
- Mumme, W.G. (1975a): Junoite,  $Pb_3Bi_8(S, Se)_6$ , a new sulfosalt from Tennant Creek, Australia: its crystal structure and relationship with other bismuth sulfosalts. *Am. Mineral.*, 60, 548-558.
- (1975b): The crystal structure of krupkaite,  $Cu_3Pb_3$  from the Juno Mine at Tennant Creek, Northern Territory, Australia. *Mineral.*, 60, 300-308.
- (1976): Proudite from Tennant Creek, Northern Territory, Australia: its crystal structure and relationship with weibullite and wittite. *Am. Mineral.*, 61, 839-852.
- (1980a): Seleniferous lead-bismuth sulphosalts from Falun, Sweden: Weibullite, wittite, and nordströmitite. *Am. Mineral.*, 65, 789-796.
- (1980b): The crystal structure of nordströmitite  $C_4Pb_4(S, Se)_4$ , from Falun, Sweden: a member of the junoite homologous series. *Can. Mineral.*, 18, 343-352.
- (1980c): Weibullite  $Ag_{0.32}Pb_{5.09}Bi_{8.55}Se_{6.08}S_{11.92}$  from Falun, Sweden: a higher homolog of galenite. *Can. Mineral.*, 18, 1-12.
- (1986): The crystal structure of perchite, a mineral of the cuprobismutite series. *Can. Mineral.*, 24, 513-521.





### 3. Other members

#### **Imhofite**, $\text{Ti}_{5.8}\text{As}_{15.4}\text{S}_{26}$

STR: Divjakovic & Nowacki (1976); Balić-Žunić & Makovicky (1993).

#### **Gillulyite**, $\text{Ti}_2\text{As}_{7.5}\text{Sb}_{0.3}\text{S}_{13}$

Def.: Wilson *et al.* (1991).

STR: Foit *et al.* (1995); Makovicky & Balić-Žunić (1999).

### 4. PbS archetype

#### **Gerstleyite**, $\text{Na}_2(\text{Sb}, \text{As})_8\text{S}_{13} \cdot 2\text{H}_2\text{O}$

STR: Nakai & Appleman (1981).

### 5.2. Rebulite plesiotypic pair

A comparative modular analysis of the crystal structures of rebulite and jankovičite has been presented by Makovicky & Balić-Žunić (1998).

#### **Rebulite**, $\text{Ti}_5\text{As}_8\text{Sb}_5\text{S}_{22}$

Def.: a complete description required for a definition is lacking.

STR: Balić-Žunić *et al.* (1982).

#### **Jankovičite**, $\text{Ti}_5\text{Sb}_9(\text{As}, \text{Sb})_4\text{S}_{22}$

Def.: Cvetkovic *et al.* (1995).

STR: Libowitzky *et al.* (1995).

### 5.3. Single type: sicherite

#### **Sicherite**, $\text{Ag}_2\text{Ti}(\text{As}, \text{Sb})_3\text{S}_6$

Definition and STR: Graeser *et al.* (2001).

### 5.4. Unclassified

#### **Erniggliite**, $\text{SnTi}_2\text{As}_2\text{S}_6$

Definition and STR: Graeser *et al.* (1992).

#### **Vrbaite**, $\text{Hg}_3\text{Ti}_4\text{As}_8\text{Sb}_2\text{S}_{20}$

STR: Ohmasa & Nowacki (1971).

#### **Simonite**, $\text{HgTiAs}_3\text{S}_6$

Def.: a complete description required for a definition is lacking.

STR: Engel *et al.* (1982).

#### **Vaughanite**, $\text{HgTiSb}_4\text{S}_7$

Def.: Harris *et al.* (1989).

#### **Gabrielite**, $\text{Cu}_2\text{AgTi}_2\text{As}_3\text{S}_7$

Def.: Graeser *et al.* (2006).

STR: Balić-Žunić *et al.* (2006). A 3-slab structure, with one of the layers related to cyclic sulfosalts.

#### **Rebulite: needs a mineralogical description**

Rebulite was defined on the basis of its crystal-structure study (Balić-Žunić *et al.*, 1982), but without a proposal submitted to the IMA-CNMNC. A complete mineralogical description is needed.

#### **Simonite: needs a mineralogical description**

Like rebulite, simonite is known only by its crystal-structure study (Engel *et al.*, 1982). It has been approved by the IMA-CNMNC, but a mineralogical description is lacking.

## 6. Sulfosalts with an excess of small (univalent) cations (Ag, Cu) relative to (As, Sb, Bi)

In the majority of these sulfosalts the ratio  $(\Sigma Me)/S$  is  $> 1$ ; however, in the presence of divalent metals (Zn, Hg, Fe), the ratio may equal 1 (galkhaite, laffittite, routhierite and stalderite).

### 6.1. Cu(Ag)-rich sulfosalts

#### 1. Wittichenite homeotypic pair

##### **Wittichenite**, $\text{Cu}_3\text{BiS}_3$

STR: Kocman & Nuffield (1973).

##### **Skinnerite**, $\text{Cu}_3\text{SbS}_3$

STR: Makovicky & Balić-Žunić (1995).

#### 2. Tetrahedrite isotypic series

Among sulfosalts, this is the most complex isotypic series, because of the multiplicity of iso- and heterovalent substitutions. Numerous crystal-structure studies have been performed since the early ones of Machatschki (1928) and Pauling & Neuman (1934). The simplified general formula is  $A_6(B, C)_6X_4Y_{12}Z$ , where  $A$  is Cu or Ag in triangular coordination,  $B$  is Cu or Ag in tetrahedral coordination,  $C$  is generally a divalent metal (typically Fe or Zn, but also Hg, Mn, Cd...) in the same tetrahedral coordination,  $X$  is Sb, As, Bi or Te in trigonal pyramidal coordination,  $Y$  is S or Se in tetrahedral coordination, and  $Z$  is S or Se in a special octahedral coordination. The presence of vacancies or interstitial atoms (*e.g.*, Cu), or heterovalent substitutions due to the incorporation of  $\text{Fe}^{3+}$  or  $\text{Te}^{4+}$ , have been confirmed by structural studies.

In tetrahedrite–tennantite, the amount of divalent metals is limited to 2 a.p.f.u. but, especially in synthetic samples, it may vary between 0 and 2, indicating the variable presence of (formal)  $\text{Cu}^{2+}$ .

There are seven well-defined species, but various data (EPMA, experimental studies, Mössbauer spectroscopy, X-ray) indicate that the crystal chemistry of this series is complex, and individual problems can require highly specialized research methods. As a consequence, limits between mineral species (for instance freibergite or goldfieldite relative to tetrahedrite) are still questionable.

Structural formulae presented below are simplified to ideal ones as much as possible (for instance, the  $B/C$  ratio is equal to  $4/2$ , and  $C$  is restricted generally to Fe and Zn). In other cases, the formula of the type sample is also given (for very rare species).

##### **Tetrahedrite**, $\text{Cu}_6[\text{Cu}_4(\text{Fe}, \text{Zn})_2]\text{Sb}_4\text{S}_{13}$

STR: Wuensch (1964); Peterson & Miller (1986); Makovicky & Skinner (1979) and Pfitzner *et al.* (1997) for Cu-pure synthetic varieties,  $\text{Cu}_{12+x}\text{Sb}_4\text{S}_{13}$ .

##### **Tennantite**, $\text{Cu}_6[\text{Cu}_4(\text{Fe}, \text{Zn})_2]\text{As}_4\text{S}_{13}$

STR: Wuensch *et al.* (1966); Makovicky *et al.* (2005) for a Cu-rich unsubstituted composition,  $\text{Cu}_{12.5}\text{As}_4\text{S}_{13}$ .

##### **Freibergite**, $\text{Ag}_6[\text{Cu}_4\text{Fe}_2]\text{Sb}_4\text{S}_{13-x} (?)$

STR: Rozhdestvenskaya *et al.* (1993).

##### **Argentotennantite**, $\text{Ag}_6[\text{Cu}_4(\text{Fe}, \text{Zn})_2]\text{As}_4\text{S}_{13}$

Type sample (Spiridonov *et al.*, 1986a):



































- (1989): The crystal structure of  $\text{Pb}_{5.05}(\text{Sb}_{3.75}\text{Bi}_{0.28})\text{S}_{10.72}\text{Se}_{0.28}$ : boulangerite of near ideal composition. *N. Jb. Miner. Mh.*, **1989** (11), 498-512.
- (1990): A note on the occurrence, composition and crystal structures of pavonite homologous series members  ${}^4\text{P}$ ,  ${}^6\text{P}$ , and  ${}^8\text{P}$ . *N. Jb. Miner. Mh.*, **1990** (5), 193-204.
- Mumme, W.G. & Watts, J.A. (1976): Pekoite,  $\text{CuPbBi}_{11}\text{S}_{18}$ , a new member of the bismuthinite–aikinite mineral series: its crystal structure and relationship with naturally- and synthetically-formed members. *Can. Mineral.*, **14**, 322-333.
- , — (1980):  $\text{HgBi}_2\text{S}_4$ : Crystal structure and relationship with the pavonite homologous series. *Acta Cryst.*, **B36**, 1300-1304.
- Mumme, W.G., Niedermayr, G., Kelly, P.R., Paar, W.H. (1983): Aschamalmite,  $\text{Pb}_{5.92}\text{Bi}_{2.06}\text{S}_9$ , from Untersulzbach Valley in Salzburg, Austria – “monoclinic heyrovskýite”. *N. Jb. Miner. Mh.*, **1983** (10), 433-444.
- Murzin, V.V., Bushmakina, A.F., Sustavov, S.G., Shcherbachev, D.K. (1996): Clerite,  $\text{MnSb}_2\text{S}_4$  – a new mineral from the Vorontsovskoye gold deposit in the Urals. *Zapiski Vsesoyuz. Mineral. Obshch.*, **124**, 95-101 (in Russian).
- Nagl, A. (1979): The crystal structure of a thallium sulfosalt,  $\text{Tl}_8\text{Pb}_4\text{Sb}_{21}\text{As}_{19}\text{S}_{68}$ . *Z. Kristallogr.*, **150**, 85-106.
- Nakai, I. & Appleman, D.E. (1981): The crystal structure of gerstleyite  $\text{Na}_2(\text{Sb,As})_8\text{S}_{13}\cdot 2\text{H}_2\text{O}$ : The first sulfosalt mineral of sodium. *Chem. Letters (Japan)*, **10**, 1327-1330.
- , — (1983): Laffittite,  $\text{AgHgAsS}_3$ : crystal structure and second occurrence from the Getchell mine, Nevada. *Am. Mineral.*, **68**, 235-244.
- Nakai, I. & Nagashima, K. (1983): crystal-chemical classification of sulfosalts – an approach towards complex inorganic compounds... in “Progress of Analytical Chemistry”, S. Fujiwara, ed. Hirokawahoten, Tokyo, 163-184 (in Japanese).
- Nakai, I., Nagashima, K., Koto, K., Morimoto, N. (1978): Crystal chemistry of oxide-chalcogenide. I. The crystal structure of sarabauite. *Acta Cryst.*, **B34**, 3569-3572.
- Nenasheva, S.N., Efimov, A.V., Sivtsov, A.V., Mozgova, N.N. (1992): Borodaevite  $[\text{Ag}_5(\text{Fe, Pb})_1\text{Bi}_7]_{13}(\text{Sb, Bi})_2\text{S}_{17}$  – a new mineral. *Zapiski Vsesoyuz. Mineral. Obshch.*, **121**, 113-120 (in Russian).
- Nespolo, M., Ferraris, G., Āurovič, S., Takéuchi, Y. (2004): Twins vs. modular crystal structures. *Z. Kristallogr.*, **219**, 773-778.
- Niizeki, N. & Buerger, M.J. (1957): The crystal structure of jamesonite,  $\text{FePb}_4\text{Sb}_6\text{S}_{14}$ . *Z. Kristallogr.*, **109**, 161-183.
- Nowacki, W. (1967): Über die mögliche Identität von “Livingit” mit Rathit-II. *N. Jb. Miner. Mh.*, **1967** (11), 353-354.
- (1968): Kristallchemie der Sulfosalze aus dem Lengenbach. *Jb. Nat. Hist. Mus. Bern*, **1966-1968**, 63-78.
- (1969): Zur Klassifikation und Kristallchemie des Sulfosalze. *Schweiz. Mineral. Petrogr. Mitt.*, **49**, 109-156.
- Nowacki, W., Itaka, Y., Burki, H., Kunz, V. (1961): Structural investigations on sulfosalts from the Lengenbach, Binn Valley (Ct. Wallis). Part 2. *Schweiz. Mineral. Petrogr. Mitt.*, **41**, 103-116.
- Nowacki, W., Marumo, F., Takeuchi, Y. (1964): Investigations on sulfides from Binntal (Canton Valais, Switzerland). *Schweiz. Mineral. Petrogr. Mitt.*, **44**, 5-9.
- Nuffield, E.W. (1975): The crystal structure of fülöppite,  $\text{Pb}_3\text{Sb}_8\text{S}_{15}$ . *Acta Cryst.*, **B31**, 151-157.
- Ohmasa, M. & Nowacki, W. (1970): A redetermination of the crystal structure of aikinite  $[\text{BiS}_2]_2[\text{Cu}^{\text{IV}}\text{Pb}^{\text{VII}}]$ . *Z. Kristallogr.*, **132**, 71-86.
- , — (1971): The crystal structure of vrbaita  $\text{Hg}_3\text{Tl}_4\text{As}_8\text{Sb}_2\text{S}_{20}$ . *Z. Kristallogr.*, **134**, 360-380.
- Ohsumi, K., Tsutsui, K., Takéuchi, Y., Tokonami, M. (1984): Reinvestigation of the lillianite structure... in “Photon Factory Activity Report 1983/1984”, VI-23.
- Orlandi, P., Dini, A., Olmi, F. (1998): Grumiplucite, a new mercury–bismuth sulfosalt species from the Levigliani mine, Apuan Alps, Tuscany, Italy. *Can. Mineral.*, **36**, 1321-1326.
- Orlandi, P., Moëlo, Y., Meerschaut, A., Palvadeau, P. (1999): Lead–antimony sulfosalts from Tuscany (Italy): I. Scainiite,  $\text{Pb}_{14}\text{Sb}_{30}\text{S}_{54}\text{O}_5$ , the first Pb–Sb oxy-sulfosalt, from Buca della Vena mine. *Eur. J. Mineral.*, **11**, 949-954.
- , —, —, — (2001): Lead–antimony sulfosalts from Tuscany (Italy). III. Pillaite,  $\text{Pb}_9\text{Sb}_{10}\text{S}_{23}\text{ClO}_{0.5}$ , a new Pb–Sb chloro-sulfosalt, from Buca della Vena mine. *Eur. J. Mineral.*, **13**, 605-610.
- Orlandi, P., Meerschaut, A., Palvadeau, P., Merlino, S. (2002): Lead–antimony sulfosalts from Tuscany (Italy): Definition and crystal structure of moëloite,  $\text{Pb}_6\text{Sb}_6\text{S}_{14}(\text{S}_3)$ , a new mineral from the Ceragiola marble quarry. *Eur. J. Mineral.*, **14**, 599-606.
- Orlandi, P., Moëlo, Y., Meerschaut, A., Palvadeau, P., Léone, P. (2004): Lead–antimony sulfosalts from Tuscany (Italy). VI. Pellouxite,  $\sim (\text{Cu, Ag})_2\text{Pb}_{21}\text{Sb}_{23}\text{S}_{55}\text{ClO}$ , a new oxy-chloro-sulfosalt, from Buca della Vena mine, Apuan Alps. *Eur. J. Mineral.*, **16**, 839-844.
- , —, —, — (2005): Lead–antimony sulfosalts from Tuscany (Italy). VIII. Rouxelite,  $\text{Cu}_2\text{HgPb}_{22}\text{Sb}_{28}\text{S}_{64}(\text{O, S})_2$ , a new sulfosalt from Buca della Vena mine, Apuan Alps: definition and crystal structure. *Can. Mineral.*, **43**, 919-933.
- Orlandi, P., Moëlo, Y., Campostrini, I., Meerschaut, A. (2007): Lead–antimony sulfosalts from Tuscany (Italy). IX. Marrucciite,  $\text{Hg}_3\text{Pb}_{16}\text{Sb}_{18}\text{S}_{46}$ , a new sulfosalt from Buca della Vena mine, Apuan Alps: Definition and crystal structure. *Eur. J. Mineral.*, **19**, 267-279.
- Otto, H.H. & Strunz, H. (1968): Zur Kristallchemie synthetischer Blei–Wismut–Spießglanze. *N. Jb. Miner. Abh.*, **108**, 1-19.
- Ozawa, T. & Nowacki, W. (1974): Note on the mineral rathite-IV. *N. Jb. Miner. Mh.*, **1974** (11), 530-531.
- Ozawa, T. & Takéuchi, Y. (1983): A new Pb–As sulfosalt, having a long periodicity, from Lengenbach. Annual Meeting Mineralogical Society Japan, 1983, Abstr., 92 (in Japanese).
- Paar, W.H., Chen, T.T., Kupčik, V., Hanke, K. (1983): Eclarit,  $(\text{Cu, Fe})\text{Pb}_9\text{Bi}_{12}\text{S}_{28}$ , ein neues Sulfosalz von Bärenbad, Hollersbachtal, Salzburg, Österreich. *Tschermaks Mineral. Petrogr. Mitt.*, **32**, 103-110.
- Paar, W.H., Putz, H., Roberts, A.C., Stanley, C.J., Culetto, F.J. (2006): Jonassonite,  $\text{Au}(\text{Bi, Pb})_5\text{S}_4$ , a new mineral species from Nagybörzsöny, Hungary. *Can. Mineral.*, **44**, 1127-1136.
- Palvadeau, P., Meerschaut, A., Orlandi, P., Moëlo, Y. (2004): Lead–antimony sulfosalts from Tuscany (Italy). VII. Crystal structure of pellouxite,  $\sim (\text{Cu, Ag})_2\text{Pb}_{21}\text{Sb}_{23}\text{S}_{55}\text{ClO}$ , an expanded monoclinic derivative of  $\text{Ba}_{12}\text{Bi}_{24}\text{S}_{48}$  hexagonal sub-type (zinkenite group). *Eur. J. Mineral.*, **16**, 845-855.
- Papp, G. (2004): History of minerals, rocks and fossil resins discovered in the Carpathian region. Hungarian Natural History Museum, Budapest, 215 p.
- Papp, G., Criddle, A.J., Stanley, C.J., Kriston, L., Nagy, G. (2008): Parajamesonite revisited: Background of the discreditation of an enigmatic mineral species. *Swiss J. Geosci.*, (online - DOI 10.1.1007/500015-007-1233-1).





